

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 199851595 B2
(10) Patent No. 729454

(54) Title
Delivery construct for antisense nucleic acids and methods of use

(51)⁷ International Patent Classification(s)
C07H 021/02 C07H 021/04
A61K 031/70 C12N 015/00

(21) Application No: 199851595

(22) Application Date: 1997.10.31

(87) WIPO No: WO98/18811

(30) Priority Data

(31) Number	(32) Date	(33) Country
08/742943	1996.10.31	US

(43) Publication Date : 1998.05.22

(43) Publication Journal Date : 1998.07.16

(44) Accepted Journal Date : 2001.02.01

(71) Applicant(s)
The Johns Hopkins University School of Medicine

(72) Inventor(s)
Harry C. Dietz

(74) Agent/Attorney
F.B. RICE and CO.,139 Rathdowne Street,CARLTON VIC 3053

(56) Related Art
PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCE USA 1996 V.93
PAGES 7219-7224
BIOCHEMISTRY 1994 VOL 33 PAGES 12127-12138

OPI DATE 22/05/98 APPLN. ID 51595/98
AOJP. DATE 16/07/98 PCT NUMBER PCT/US97/19849



AU9851595

PCT)

(51) International Patent Classification ⁶ : C07H 21/02, 21/04, A61K 31/70, C12N 15/00		AI	(11) International Publication Number: WO 98/18811
			(43) International Publication Date: 7 May 1998 (07.05.98)
(21) International Application Number: PCT/US97/19849		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BI, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 31 October 1997 (31.10.97)		Published With international search report.	
(30) Priority Data: 08/742,943 31 October 1996 (31.10.96) US			
(71) Applicant: THE JOHNS HOPKINS UNIVERSITY SCHOOL OF MEDICINE [US/US]; 720 Rutland Avenue, Baltimore, MD 21205 (US).			
(72) Inventor: DIETZ, Harry, C.; 1104 Ryegate Road, Towson, MD 21286 (US).			
(74) Agent: HAILE, Lisa, A.; Fish & Richardson P.C., Suite 1400, 4225 Executive Square, La Jolla, CA 92037 (US).			
(54) Title: DELIVERY CONSTRUCT FOR ANTISENSE NUCLEIC ACIDS AND METHODS OF USE			
(57) Abstract <p>A novel nucleic acid construct for delivery of antisense targeting sequences is provided. The construct includes intact stem loop structures and an antisense nucleic acid. Optionally, a ribozyme nucleic acid is included in the construct. The construct is useful for inhibition of selected genes in a cell. This allele-specific targeting is also useful in combination with replacement gene therapy.</p>			

NUC 37330

**DELIVERY CONSTRUCT FOR ANTISENSE NUCLEIC ACIDS
AND METHODS OF USE**

5 **Field of the Invention**

This invention relates generally to the field of delivery vehicles for nucleic acid molecules and specifically to a novel construct for delivery of antisense targeting sequences to a cell.

10 **Background of the Invention**

The use of antisense oligonucleotides offers advantages over other therapeutic regimes due to their potential for target specificity. For example, conventional chemotherapy for neoplastic and virus-related diseases has the disadvantage of systemic toxicity. The therapeutic index for chemotherapeutic agents is relatively narrow, since such agents are unable to distinguish between normal and diseased cells. Antisense oligonucleotides have the potential to be many orders of magnitude more specific than traditional drugs due to their greater number of interactions with a particular target site. In theory, an oligonucleotide of more than 15-17 nucleotides in length could have the base pairing specificity to interact with only one target gene within the entire human genome. Thus, antisense oligonucleotides have the potential specificity that could serve as a powerful tool for the study of specific gene function and as therapeutic agents for disease-causing genes.

In contrast to drugs, antisense molecules are relatively simple to design.

25 The interaction between an antisense oligonucleotide and a target mRNA is governed primarily by the sequence of the target. Oligonucleotides targeting the start codon and extending upstream or downstream have been shown to be effective. Similarly, oligonucleotides that are complementary to the splice sites have proved effective.

- 2 -

Antisense technologies for the targeted inhibition of gene expression could provide an effective strategy for the management of inherited disorders with dominant-negative or gain-of-function pathogenetic mechanisms, for the suppression of oncogenes, or for the control of a variety of infectious agents. Pathologic disorders
5 that are currently targeted by antisense therapeutics include viral infections, inflammatory disorders, cardiovascular disease, cancers, genetic disorders and autoimmune diseases. Synthetic oligodeoxynucleotides (ODNs), especially phosphorothiates and methylphosphonates, offer the advantage of enhanced stability in biological fluids and an effectively limitless supply.

10 Antisense oligonucleotides are also useful for the production of transgenic animals having alterations at the germline level, such as knockout mutations, which can be used for the study of new genes or the study of the function of a known gene. Further, antisense technology combined with gene therapy is useful for example, for suppression of expression of a mutant gene product. Such gene therapy would be
15 most advantageous in combination with a replacement regimen utilizing the "normal" gene to provide a "normal" gene product.

Unfortunately, the effective use of antisense oligonucleotides has been limited due to several problems. Disadvantages include the transient nature of ODNs, and their toxicity and propensity for producing non-sequence specified
20 biological effects. Other disadvantages include low expression or limited stability of complementary RNAs which result in their nonspecific targeting or low efficiency of target inhibition. Antisense oligonucleotides are often poorly taken up by cells and therefore may never reach their target site. Often, antisense oligonucleotides do not reach the nucleus of a cell once administered, the site of their RNA and DNA targets.
25 In certain applications the antisense molecules are microinjected directly into the cells. This technique works well in the laboratory, however, it cannot be applied to patients. Many of the studies with antisense show that gene expression is suppressed by 80-90% of the normal level, however, such reduction is not typically sufficient to reduce the biological effect, e.g., 10-20% expression is sufficient to maintain the
30 biological function sought to suppress.

- 3 -

There is a need to develop a delivery system for antisense molecules that gives the antisense enhanced stability, for example by being resistant to nuclease activity or by being enriched in the nucleus, while still allowing specificity of the antisense for its target RNA or DNA. Such a system would provide effective

5 targeting of the message with the end result being significant inhibition of expression of a particular gene.

- 4 -

Summary of the Invention

The present invention provides a novel nucleic acid construct for delivery of antisense targeting sequences for inhibition of selected genes in a cell. The construct includes intact stem loop structures and an antisense nucleic acid. This
5 allele-specific targeting is also useful in combination with replacement gene therapy.

In a first embodiment, the invention provides a nucleic acid construct for suppressing gene expression. The construct includes a 5' stem loop structure, an antisense nucleic acid, and a 3' stem loop structure. Such "unmodified" or "intact" stem loop structures flank the antisense nucleic acid so that the antisense
10 oligonucleotides can readily interact with any target sequence. In a preferred embodiment, the stem loop structures are U snRNA stem loops, and most preferably, U1 snRNA stem loops. The construct provides a cloning site outside of the stem loop structures, thus rendering the stem loop structures "unmodified," into which virtually any antisense oligonucleotide could be inserted. In contrast to previous antisense
15 delivery vehicles which have been designed to interfere with mRNA splicing, the construct of the present invention does not necessarily affect the splicing machinery so as not to disrupt normal cellular mRNA processing. In other words, while previous constructs retain elements that interact with spliceosome proteins, the construct of the invention eliminates such elements. Preferably, a U1 snRNA promoter is included in
20 the construct. Optionally, a ribozyme nucleic acid is included in the construct.

In another embodiment, the invention provides a method for suppression of gene expression whereby a suppressive-effective amount of the nucleic acid construct of the invention is administered to a cell thereby suppressing expression of the gene. The method is particularly useful when used in combination with
25 replacement gene therapy which provides a gene which encodes and can reconstitute the wild-type protein and which is resistant to targeting by the antisense and/or ribozyme.

The seminal discovery of the delivery vehicle of the invention now allows transgenic knockout animals to be created for the study of known or as of yet

unknown genes. For example, phenotypic consequences of loss of function due to targeted disruption of a gene(s) can be determined.

- 5 Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215

- 6 -

Brief Description of the Drawings

Figure 1 shows an illustrative expression construct of the invention.

Figure 2 shows the sequence and predicted structure of chimeric transcripts derived from pU1/FIB.

5 Figure 3 shows an immunohistochemical analysis of MG63 cells that were stably transfected with an expression construct (pZeoSVLacZ, Invitrogen) lacking targeting sequence (panels A and C) or pU1/FIB (panels B and D).

Figure 4 shows a Northern blot analysis of 3.5 µg of poly(a) RNA extracted from confluent MG63 cells that were untransfected (lane 1), or stably
10 transfected with either pZeoSVLacZ (Lane 2) or pU1/FIB (lane 3-5).

- 7 -

Description of the Preferred Embodiments

The present invention provides a novel nucleic acid construct which provides a general means of delivery for antisense targeting sequences. The construct includes stabilizing structural elements, stem loop structures, which increase the effectiveness of the targeting construct. Also included is a method of suppression of gene expression utilizing the novel construct of the invention.

The nucleic acid construct of the invention which is useful for suppressing gene expression includes 5' and 3' stem loop structures and an antisense nucleic acid. While not wanting to be bound by a particular theory, it is believed that the stem loop structures provide enhanced stability of the targeting molecules by conferring resistance to exonucleases and also promote duplex formation and stability.

The "stem loop" structures refer to nucleic acid structures that have folding patterns which form hairpins and flank the antisense targeting sequence. The stem loop structures are preferably unmodified, naturally occurring structures. Alternatively, one of skill in the art would be able to synthesize such structures to "mimic" the naturally occurring structures. The preferred stem loop structures in the construct of the invention are unmodified U snRNA structures. The term "unmodified" means that the folding pattern of the stem loop structure is not compromised by alterations in the nucleic acid sequence of the naturally occurring molecule. For example, it is understood that alterations which include, but are not limited to, mutations, insertions, deletions and substitutions of one or more nucleotides can be made within the sequence of the stem loop, as long as the stabilization function and hairpin formation of the stem loop is maintained.

Preferably, the stem loop structures are derived from or synthesized according to the nucleotide sequence of the U snRNA stem loop structures. An exemplary construct utilizing U1 snRNA stem loops is shown in FIGURE 1 herein. One can produce a chimeric construct having a 5' stem loop structure from one species and a 3' structure from a second species. For example, a 5' stem loop is derived from human U snRNA and the 3' stem loop derived from a yeast U snRNA.

- 8 -

Other snRNA stem loop structures can be derived from other mammals (e.g., porcine, bovine), or yeast for example. Other chimeric constructs can be prepared using a 5' stem loop structure from one snRNA (e.g., U1) and a 3' stem loop structure from another snRNA (e.g., U3). Virtually any combination of stem loop structures is
5 envisioned in the construct of the invention.

As used herein, the term "nucleic acid" or nucleic acid sequence" refers to a polymer of deoxyribonucleotides or ribonucleotides, in the form of a separate fragment or as a component of a larger construct. For example, nucleic acids can be assembled from cDNA fragments or from oligonucleotides to generate a synthetic
10 gene which is capable of being expressed in a recombinant transcriptional unit. Polynucleotide or nucleic acid sequences of the invention include DNA, RNA and cDNA sequences.

Nucleic acid sequences utilized in the invention can be obtained by several methods. Sequences for specific genes or stem loop structures can determined from
15 published sequences and can also be found in GenBank, National Institutes of Health computer database. Nucleic acids can then be chemically synthesized by standard methods for example.

The construct of the invention includes an antisense nucleic acid flanked by the stem loop structures. The antisense nucleic acid can be directed toward any
20 target nucleic acid, and preferably to a target message (mRNA). Any antisense which includes sequences capable of hybridizing with its complementary target can be used in the construct of the invention. For example, antisense sequences can be directed to the 5' or 3' termini of the target message, to splice junctions, or to internal sequences. One of skill in the art will readily be able to determine which sequences
25 to use as the appropriate antisense nucleic acid construct. "Antisense nucleic acids" are DNA or RNA molecules that are complementary to at least a portion of a specific mRNA molecule (Weintraub, Scientific American, 262:40, 1990). In the cell, the antisense nucleic acids hybridize to the corresponding mRNA, forming a double-stranded molecule. The antisense nucleic acids interfere with the translation of the
30 mRNA, since the cell will not translate a mRNA that is double-stranded. Antisense

- 9 -

oligomers of about 15 nucleotides are preferred, since they are easily synthesized and are less likely to be toxic than larger molecules when introduced into the target cell.

The use of antisense methods to inhibit the *in vitro* translation of genes is well known in the art (Marcus-Sakura, *Anal. Biochem.*, 172:289, 1988). An antisense core nucleic

5 acid may contain about 10 nucleotides complementary to the target message.

Examples of target messages include transcription regulatory factors (*e.g.*, *rent-1*); viral encoded proteins (*e.g.*, human papilloma virus E6, human immunodeficiency virus tat), (*e.g.*, haluronic acid synthase), structural proteins (*e.g.*, fibrillin), cytokines, oncogenes and growth factors (*e.g.*, interleukins), etc. It should be noted that the

10 antisense nucleic acid can be useful for reducing the expression of either normal or aberrant genes.

The antisense nucleic acid can be used to block expression of a mutant protein or a dominantly active gene product, such as amyloid precursor protein that accumulates in Alzheimer's disease. Such methods are also useful for the treatment of

15 Huntington's disease, hereditary Parkinsonism, and other diseases. Antisense nucleic acids are also useful for the inhibition of expression of proteins associated with toxicity or gene products introduced into the cell, such as those introduced by an infectious agent (*e.g.*, a virus).

It may be desirable to transfer an antisense nucleic acid encoding a

20 biological response modifier in order to reduce the expression of such biological response modifier. Included in this category are nucleic acids encoding immunopotentiating agents including a number of the cytokines classified as "interleukins". These include, for example, interleukins 1 through 12. Also included in this category, although not necessarily working according to the same mechanisms,

25 are interferons, and in particular gamma interferon (γ -IFN), tumor necrosis factor (TNF) and granulocyte-macrophage-colony stimulating factor (GM-CSF). It may be desirable to deliver such antisense nucleic acids to cells of the immune system to treat enzymatic related disorders or immune defects. Antisense nucleic acids can be used to reduce expression of growth factors, toxic peptides, ligands, receptors, or other

30 physiologically important proteins.

- 10 -

The construct of the invention may further include a ribozyme nucleic acid. Ribozymes are RNA molecules possessing the ability to specifically cleave other single-stranded RNA. Through the modification of nucleotide sequences which encode these RNAs, it is possible to engineer molecules that recognize specific
5 nucleotide sequences in an RNA molecule and cleave it (Cech, *J. Amer. Med. Assn.*, 260:3030, 1988). A major advantage of this approach is that, because the ribozymes are engineered to be sequence-specific, only mRNAs with sequences complementary to the construct containing the ribozyme are inactivated.

There are two basic types of ribozymes namely, tetrahymena-type
10 (Hasselhoff, et al., *Nature*, 334:585, 1988) and "hammerhead"-type. Tetrahymena-type ribozymes recognize sequences which are four bases in length, while "hammerhead"-type ribozymes recognize base sequences from about 3 to 18 bases in length. The longer the recognition sequence, the greater the likelihood that the sequence will occur exclusively in the target mRNA species. Consequently,
15 hammerhead-type ribozymes are preferable to tetrahymena-type ribozymes for inactivating a specific mRNA species. The preferred ribozyme of the present invention is a hammerhead type ribozyme. The target message is required to contain a ribozyme cleavage site sequence such as 5'-GUC-3' or 5'-GUA-3' when a ribozyme is included in the construct.

20 The construct may also optionally include a triplex oligomer. Use of an oligonucleotide to stall transcription is known as the triplex strategy since the oligomer winds around double-helical DNA, forming a three-strand helix. Therefore, these triplex compounds can be designed to recognize a unique site on a chosen gene (Maher, et al., *Antisense Res. and Dev.*, 1(3):227, 1991; Helene, C., *Anticancer Drug*
25 *Design*, 6(6):569, 1991).

The phrase "nucleic acid sequence expressing a product of interest" refers to a nucleic acid sequence which when expressed results in a product selected from a protein or antisense RNA, for example. The term "structural gene" excludes the non-coding regulatory sequence which drives transcription. The structural gene may
30 be derived in whole or in part from any source known to the art, including a plant, a

- 11 -

fungus, an animal, a bacterial genome or episome, eukaryotic, nuclear or plasmid DNA, cDNA, viral DNA or chemically synthesized DNA. A structural gene may contain one or more modifications in either the coding or the untranslated regions which could affect the biological activity or the chemical structure of the expression product, the rate of expression or the manner of expression control. Such modifications include, but are not limited to, mutations, insertions, deletions and substitutions of one or more nucleotides. The structural gene may constitute an uninterrupted coding sequence or it may include one or more introns, bound by the appropriate splice junctions.

10 The term "operably associated" refers to functional linkage between the regulatory (e.g., promoter) sequence and the nucleic acid regulated by the regulatory sequence. The operably linked regulatory sequence controls the expression of the product expressed by the structural gene or antisense. The regulatory sequence may be homologous or heterologous to the desired gene sequence. A wide range of
15 promoters may be utilized, including viral or mammalian promoters. Cell- or tissue-specific promoters can be utilized to target expression of gene sequences in specific cell populations. Suitable mammalian and viral promoters for the present invention are available in the art. A preferred promoter in the construct of the invention is a U snRNA promoter, and most preferably, a U1 snRNA promoter.

20 The choice of a particular heterologous promoter region as a "regulatory nucleotide sequence" of the invention, is dictated by the spatial and temporal pattern of expression that is desired for the transactivator gene and ultimately for the target transgene. Promoter regions of the invention include eukaryotically derived promoters which predominantly direct expression in, for example, the reproductive
25 system (e.g., breast, ovary, testes); the musculoskeletal system (e.g., muscle or joint tissue); the cardiovascular system (e.g., capillaries or heart); the respiratory system (e.g., lung or nasal passages); the urological system (e.g., kidney or bladder); the gastrointestinal system (e.g., pancreas, liver, or intestines); the immune system (e.g., thymus, spleen, or circulating immunological cells); the endocrine system (e.g.,
30 pituitary, gonads, and thyroid); the nervous system (e.g., neurons); and the

- 12 -

hematopoietic system (e.g., bone marrow and peripheral blood). Further, promoters of the invention include, but are not limited to the elastase promoter (including its enhancer; expression in pancreatic acinar cells); the alpha-A-crystallin promoter (expression in the eye lens tissue); the insulin promoter (including enhancer; expression in the pancreatic beta cells) and the albumin promoter region, including its enhancer.

Alternatively, non-eukaryotically-derived promoters such as virally-derived and prokaryotically-derived promoters are also included in the present invention. Such virally-derived promoters include, but are not limited to MMTV and MoSV LTR, SV40 early region, RSV or CMV LTR, which direct expression of viral or host genes in specific tissues and in many cell types.

Promoters useful in the invention include both constitutive and inducible natural promoters as well as engineered promoters. To be most useful, an inducible promoter should 1) provide low expression in the absence of the inducer; 2) provide high expression in the presence of the inducer; 3) use an induction scheme that does not interfere with the normal physiology of the cell; and 4) have no effect on the expression of other genes. Both constitutive and inducible promoters and enhancers will be known to those of skill in the art. The promoters used in the construct of the present invention may be modified, if desired, to affect their control characteristics.

The regulatory nucleotide sequence of the invention may also include "enhancer" regions. Enhancers, as used herein, refer to DNA sequences which affect transcription of a gene by RNA polymerase II, without regard to position or orientation. An enhancer region can be thousands of base pairs from a transcription unit and still affect its transcription. Example of enhancers known in the art include the immunoglobulin heavy (mu) chain or light (kappa) chain enhancers and the SV40 72-base pair repeats.

It may be desirable to deliver the construct of the invention directly to the cell without the requirement for expression, therefore, a promoter or other regulatory nucleic acid is optional. One example of a targeted delivery system for antisense polynucleotides is a colloidal dispersion system. Colloidal dispersion systems include

- 13 -

macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes or liposome formulations. The preferred colloidal system of this invention is a liposome. Liposomes are artificial membrane vesicles which are useful as delivery vehicles *in vitro* and *in vivo*. These formulations may have net cationic, anionic or neutral charge characteristics are useful characteristics with *in vitro*, *in vivo* and *ex vivo* delivery methods. It has been shown that large unilamellar vesicles (LUV), which range in size from 0.2-4.0 μm can encapsulate a substantial percentage of an aqueous buffer containing large macromolecules. RNA, DNA and intact virions can be encapsulated within the aqueous interior and be delivered to cells in a biologically active form (Fraley, *et al.*, *Trends Biochem. Sci.*, 6:77, 1981). In addition to mammalian cells, liposomes have been used for delivery of polynucleotides in plant, yeast and bacterial cells. In order for a liposome to be an efficient gene transfer vehicle, the following characteristics should be present: (1) encapsulation of the genes of interest at high efficiency while not compromising their biological activity; (2) preferential and substantial binding to a target cell in comparison to non-target cells; (3) delivery of the aqueous contents of the vesicle to the target cell cytoplasm at high efficiency; and (4) accurate and effective expression of genetic information (Mannino, *et al.*, *Biotechniques*, 6:682, 1988).

The composition of the liposome is usually a combination of phospholipids, particularly high-phase-transition-temperature phospholipids, usually in combination with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical characteristics of liposomes depend on pH, ionic strength, and the presence of divalent cations.

The construct of the invention can also be delivered as a naked "gene expression vector". This means that the construct is not associated with a delivery vehicle (e.g., liposomes, colloidal particles and the like). One of the principal advantages touted for naked DNA vectors has been the lack of immune responses stimulated by the vector itself.

- 14 -

The backbone or framework of the construct preferably includes U1 snRNA nucleic acid sequences since U1 snRNA is enriched in the nucleus, is widely dispersed in the nucleoplasm, and is abundantly expressed. In the exemplary construct shown in FIGURE 1, an "antisense targeting core" was substituted for the

5 Sm protein binding site between the two naturally occurring U1 snRNA hairpins. This core contains a 30-35 basepair sequence that is directly complementary to the target message, interrupted in its center by an autocatalytic hammerhead ribozyme loop. Duplex formation aligns the ribozyme with the GUC or GUA consensus sites for ribozyme cleavage within the target message. An illustrative expression construct

10 of the invention which targets RENT1 transcripts, which are involved in the NMRD pathway, described below, is shown in Figure 1. The term *rent1* and RENT1 (regulator of nonsense transcripts) refers to the gene and protein, respectively, from either the murine or human species.

The construct described above is useful for suppressing gene expression.

15 For example, it may be desirable to modulate the expression of a gene when it is over-expressed. Where a cell proliferative disorder is associated with the expression of a gene, nucleic acid sequences that interfere with the gene's expression at the translational level can be used. This approach utilizes, for example, antisense nucleic acid, ribozymes, or triplex agents to block transcription or translation of a specific

20 mRNA, either by masking that mRNA with an antisense nucleic acid or triplex agent, or by cleaving it with a ribozyme, as described above.

The construct may also optionally include a 5' cap structure, such as a trimethylguanosine cap.

In another embodiment, the present invention also includes a method for

25 suppression of gene expression which includes administering to a cell a suppressive-effective amount of the nucleic acid construct of the invention so that the expression of the gene is suppressed. The term "suppressive-effective" amount means that amount of the construct, and thus antisense, administered is sufficient to suppress the expression of the target, *e.g.*, inhibit translation of mRNA, by at least 75% of the

30 normal expression, and preferably by at least 90%. The effectiveness of the construct

- 15 -

can be determined phenotypically or by standard Northern blot analysis or immunohistochemically, for example. Other standard nucleic acid detection techniques or alternatively immunodiagnostic techniques will be known to those of skill in the art (e.g., Western or Northwestern blot analysis).

5 The present invention also provides a method of gene therapy for the treatment of cell proliferative or immunologic disorders and diseases such as which are mediated by various proteins. The term "cell-proliferative disorder" denotes malignant as well as non-malignant cell populations which often appear to differ from the surrounding tissue both morphologically and genotypically. Such disorders may
10 be associated, for example, with abnormal expression of a gene. "Abnormal expression" encompasses both increased or decreased levels of expression, as well as expression of a mutant form of a gene such that the normal function of the gene product is altered. Abnormal expression also includes inappropriate expression of during the cell cycle or in an incorrect cell type. The antisense polynucleotide is
15 useful in treating malignancies of the various organ systems. Such therapy would achieve its therapeutic effect by introduction of the antisense construct into cells having the proliferative disorder. Delivery of antisense polynucleotide, can be achieved using a recombinant expression vector such as a chimeric virus or a colloidal dispersion system as described above.

20 The construct may also be useful in treating malignancies of the various organ systems, such as, for example, lung, breast, lymphoid, gastrointestinal, and genito-urinary tract as well as adenocarcinomas which include malignancies such as most colon cancers, renal-cell carcinoma, prostate cancer, leukemia, breast cancer, non-small cell carcinoma of the lung, cancer of the small intestine, and cancer of the
25 esophagus.

 The method is also useful in treating non-malignant or immunologically-related cell-proliferative diseases such as psoriasis, pemphigus vulgaris, Bechet's syndrome, acute respiratory distress syndrome (ARDS), ischemic heart disease, post-dialysis syndrome, rheumatoid arthritis, acquired immune deficiency syndrome,
30 vasculitis, lipid histiocytosis, septic shock and inflammation in general.

- 16 -

Various viral vectors which can be utilized for gene therapy as taught herein include adenovirus, adeno-associated virus, herpes virus, vaccinia, or, an RNA virus such as a retrovirus. Preferably, the retroviral vector is a derivative of a murine or avian retrovirus. Examples of retroviral vectors in which a single foreign gene can be inserted include, but are not limited to: Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), and Rous Sarcoma Virus (RSV). Preferably, when the subject is a human, a vector such as the gibbon ape leukemia virus (GaLV) is utilized. A number of additional retroviral vectors can incorporate multiple genes. All of these vectors can transfer or incorporate a gene for a selectable marker so that transduced cells can be identified and generated. By inserting a sequence of interest into the viral vector, along with another gene which encodes the ligand for a receptor on a specific target cell, for example, the vector is now target specific. Retroviral vectors can be made target specific by attaching, for example, a sugar, a glycolipid, or a protein. Preferred targeting is accomplished by using an antibody to target the retroviral vector. Those of skill in the art will know of, or can readily ascertain without undue experimentation, specific polynucleotide sequences which can be inserted into the retroviral genome or attached to a viral envelope to allow target specific delivery of the retroviral vector containing the antisense polynucleotide.

Administration of the construct of the invention can be *in vivo*, *in vitro* or *ex vivo*.

The method of the invention also envisions gene replacement therapy to reconstitute expression of a wild-type protein in cells expressing a defective protein or, in the case of a heterozygote, a defective and a "normal" protein. Therefore, in another embodiment the invention provides a method of suppressing gene expression as above and further comprises administering a modified nucleic acid encoding a wild-type polypeptide corresponding to the gene product of the gene being suppressed, wherein the modified nucleic acid is resistant to ribozyme cleavage and/or antisense inhibition. Due to degeneracy in the genetic code, multiple nucleotide substitutions can be made across the antisense oligonucleotide target

- 17 -

region of the coding sequence within an exogenously supplied expression construct encoding the wild-type gene of interest. Such substitutions would inhibit duplex formation between the gene suppressed and the antisense nucleic acid and transcripts derived from the replacement "normal" gene, while maintaining the fidelity of the amino acid sequence. In other words, the replacement gene would provide a protein with the correct amino acid sequence utilizing codons with nucleotides that are not recognized by the antisense construct and/or ribozyme. In this method, it is therefore possible to "knockout" a defective copy of a gene/mRNA and introduce a "normal" copy resistant to the "knockout" construct. Thus, both normal and aberrant expression of a gene can be blocked by introduction of the antisense construct of the invention; to restore normal function of the gene, a copy of the wild-type gene is introduced, that, while coding for a normal protein, is resistant to duplex formation with the antisense construct and/or ribozyme cleavage.

There are a number of inherited diseases in which defective genes may be replaced including: lysosomal storage diseases such as those involving β -hexosaminidase or glucocerebrosidase; deficiencies in hypoxanthine phosphoribosyl transferase activity (the "Lesch-Nyhan" syndrome); amyloid polyneuropathies (-prealbumin); Duchenne's muscular dystrophy, and retinoblastoma, for example.

Pathologic disorders that can be targeted using the method of the invention include but are not limited to viral infections, inflammatory disorders, cardiovascular disease, cancers, genetic disorders and autoimmune diseases.

The method of the present invention is also useful for suppression of a dominant negative mutation. The term "dominant negative" is a term of art and refers to expression of a gene resulting in a gene product that actively interferes with the function of a "normal", endogenous protein. Thus, a mutant protein or dominantly active gene product, such as amyloid precursor protein that accumulates in Alzheimer's disease can be blocked. The dominant negative phenotype is conveyed by the expression of the mutant protein that interferes with the function of the normal protein. Such an effect is similar to the effect of dominance of one allele of a pair of alleles encoding homologous genes on a pair of homologous chromosomes such that

- 18 -

the phenotypic effect of the one allele exerts a deleterious controlling influence over the other allele.

The nucleic acid construct can be utilized in the discovery of unknown human genes, for example, with *a priori* knowledge of the gene in question. In one embodiment of this invention, cells are isolated from an individual displaying a particular phenotype which is suspected of being caused by a gene or genes containing nonsense mutations. A subset of these cells is contacted (e.g., by transfecting the cells), with the construct that contains at least an antisense nucleic acid for suppression or inhibition of the expression or function of a gene (e.g., Rent-1 gene or other genes involved in NMRD) or its gene product in the cells. Levels of mRNA containing nonsense mutations in these cells are elevated to higher levels. Using standard techniques of differential display of mRNA, or genetic subtraction techniques commonly applied to cDNAs, the mRNA containing nonsense mutations can be enriched and then isolated by comparing the mRNA populations of the original cells with the mRNA populations in cells containing the antisense.

In a further embodiment, a transgenic animal can be developed using the novel construct and method of the invention in order to identify the impact of increased or decreased gene expression on a particular pathway or phenotype. Protocols useful in producing such transgenic animals are described below. The protocol generally follows conventional techniques for introduction of expressible transgenes into mammals. Those of ordinary skill in the art will be familiar with these applications and will be able to apply the techniques in the context of the present invention without undue experimentation.

For example, embryonic target cells at various developmental stages can be used to introduce transgenes. Different methods are used depending on the stage of development of the embryonic target cell. The zygote is the best target for microinjection. In the mouse, the male pronucleus reaches the size of approximately 20 micrometers in diameter which allows reproducible injection of 1-2pl of DNA solution. The use of zygotes as a target for gene transfer has a major advantage in that in most cases the injected DNA will be incorporated into the host gene before the

- 19 -

first cleavage (Brinster, *et al.*, *Proc. Natl. Acad. Sci. USA* 82:4438-4442, 1985). As a consequence, all cells of the transgenic non-human animal will carry the incorporated transgene. In general, this will also be reflected in the efficient transmission of the transgene to offspring of the founder since 50% of the germ cells will harbor the transgene. Microinjection of zygotes is a preferred method for incorporating transgenes in practicing the invention.

Retroviral infection can also be used to introduce transgene into a non-human animal. The developing non-human embryo can be cultured *in vitro* to the blastocyst stage. During this time, the blastomeres can be targets for retroviral infection (Jaenisch, *Proc. Natl. Acad. Sci. USA* 73:1260-1264, 1976). Efficient infection of the blastomeres is obtained by enzymatic treatment to remove the zona pellucida (Hogan, *et al.*, *Manipulating the Mouse Embryo*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). The viral vector system used to introduce the transgene is typically a replication-defective retrovirus carrying the transgene (Jahner, *et al.*, *Proc. Natl. Acad. Sci. USA*, 82:6927-6931, 1985; Van der Putten, *et al.*, *Proc. Natl. Acad. Sci. USA* 82:6148-6152). Transfection is easily and efficiently obtained by culturing the blastomeres on a monolayer of virus-producing cells (Van der Putten, *supra*; Steward, *et al.*, *EMBO J.*, 6:383-388, 1987).

Alternatively, infection can be performed at a later stage. Virus or virus-producing cells can be injected into the blastocoele (Jahner, *et al.*, *Nature*, 298:623-628, 1982). Most of the founders will be mosaic for the transgene since incorporation occurs only in a subset of the cells which formed the transgenic non-human animal. Further, the founder may contain various retroviral insertions of the transgene at different positions in the genome which generally will segregate in the offspring. In addition, it is also possible to introduce transgenes into the germ line, albeit with low efficiency, by intrauterine retroviral infection of the midgestation embryo (Jahner, *et al.*, *supra*, 1982).

A third type of target cell for introduction of heterologous nucleic acid sequences is the embryonal stem cell (ES). ES cells are obtained from pre-implantation embryos cultured *in vitro* and fused with embryos (Evans, *et al.*, *Nature*,

- 20 -

292:154-156, 1981; Bradley, *et al.*, *Nature*, 309:255-258, 1984; Gossler, *et al.*, *Proc. Natl. Acad. Sci USA*, 83:9065-9069, 1986; and Robertson, *et al.*, *Nature*, 322:445-448, 1986). Transgenes can be efficiently introduced into the ES cells by DNA transfection or by retrovirus-mediated transduction. These transformed ES cells can

5 thereafter be combined with blastocysts from a non-human animal. The ES cells will thereafter colonize the embryo and contribute to the germ line of the resulting chimeric animal (see for review, Jaenisch, *Science*, 240:1468-1474, 1988). Any ES cell may be used in accordance with the present invention. It is, however, preferred to use primary isolates of ES cells. Such isolates may be obtained directly from

10 embryos such as with the CCE cell line disclosed by Robertson, E.J., in *Current Communications in Molecular biology*, Capecchi, M.R. (Ed.) Cold Springs Harbor Press, Cold Springs Harbor, NY (1989), pp.39-44), or from the clonal isolation of ES cells from the CCE cell line (Schwartzberg, P.A. *et al.*, *Science* 246:799 (1989). ES cells may be derived or isolated from any species, although cells derived or isolated

15 from mammals such as rodents, rabbits, and non-human primates are preferred.

Construction of transgenes can be performed by those of ordinary skill in the art using the teachings herein. One of ordinary skill in the art can "knock out" a gene in mice by targeted disruption of the gene. This can be accomplished by homologous recombination in murine embryonic stem (ES) cells using standard

20 techniques. The clinical and cellular consequences of targeted disruption can be investigated in multiple genetic backgrounds including inbred strains, strains with many undefined nonsense alleles, and strains of known mutant genotype to determine if: (a) targeted disruption of the gene can effect a loss of function; (b) loss of function can have an phenotype consequences (i.e., the creation of knockout phenotypes), (c)

25 loss of the gene product unmasks the effects of protein that can be expressed from "physiologic" somatically acquired or inherited nonsense alleles upon transcript stabilization.

The cre/lox system as described in US Patent Serial No. 4,959,317, incorporated herein by reference, can be utilized in the production of transgenic

30 animals. A first and second loxP DNA sequence is introduced into cells connected by

- 21 -

a pre-selected antisense or replacement gene, herein referred to as a "target transgene". The "target transgene" of interest can be a complete gene or any other sequence of nucleotides including those of homologous, heterologous, or synthetic origin. The target transgene sequence can be for example, an antisense or
5 replacement gene for a structural protein, an enzyme, or a regulatory molecule. The target transgene may also be a gene of undetermined function. Using tissue-specific or developmentally-specific regulatory sequences (as described above) to direct expression of the target transgene, a function could be identified. If the first and second lox sites have the same orientation (direct repeats), activation of the regulatory
10 nucleotide sequence of the transactivator transgene results in a deletion of the target transgene DNA, such that ablation or modification of activity results. If the first and second lox sites have opposite orientation (inverted repeats), activation of the regulatory nucleotide produces an inversion of the nucleotides sequence of the target transgene.

15 The construct of the invention may be used to introduce DNA sequences into the germ line cells of "non-humans" to create transgenic animals. The preferred animal of the invention is a mouse. However, other non-humans of the invention include but are not limited to other rodents (e.g. rat, hamster), rabbits, chickens, sheep, goats, fish, pigs, cattle, and non-human primates.

20 In yet another embodiment, an antisense library can be constructed using the nucleic acid construct of the invention in order to identify novel genes. For example, a U1 construct library containing antisense core sequences (e.g., random sequences or sequences derived from a cDNA library) is transfected into host cells and clones exhibiting a desired phenotype are identified. Such clones are then
25 examined to determine the nature of the regulatory sequence in the construct.

Therefore, novel genes having specific functions can be identified. For example, cells that acquire a transformed phenotype may contain an antisense molecule for a novel tumor suppressor gene. A match between a given suppressive nucleic acid and a previously cloned gene can be made using standard techniques (e.g., BLAST search
30 algorithm).

- 22 -

The following Examples are intended to illustrate, but not to limit the invention. While such Examples are typical of those that might be used, other procedures known to those skilled in the art may alternatively be utilized.

EXAMPLES

The following examples provide a description of construction of an exemplary nucleic acid construct of the invention, utilizing 5' and 3' unmodified stem loop structures from U1 snRNA which flank an antisense molecule directed toward the fibrillin-1 gene (mRNA).

EXAMPLE 1

An antisense expression construct was constructed that incorporates several potentially enhancing features. The pU1/FIB vector was constructed on the backbone of the pZeoSV (Invitrogen) prokaryotic/eukaryotic expression vector. The SV40 promoter, polyadenylation site and polylinker were excised from pZeoSV at the BamHI sites. A U1 snRNA expression cassette cloned into pUC13 was excised with BamHI digestion and ligated into the BamHI sites of the modified pZeoSV. Two rounds of site-directed mutagenesis (Deng, et al., *Anal. Biochem.*, 200, 81; 1992) were then performed to change four nucleotides flanking the Sm protein restriction sites (pZeoU1EcoSpe). Complementary oligonucleotides that encode the antisense 'core' sequence, shown in Figure 1, including the 24 highly conserved nucleotides of hammerhead ribozymes (Cech, et al., *Ann. Rev. Biochem.*, 55:599, were synthesized and annealed at 40° C such that the remaining 5' and 3' overhangs were exactly complementary to the overhangs left by EcoRI and SpeI digestion. The sequences of the oligonucleotides were as follows:

5'-AATTGGCGATCTCCAGCACTGATGAGTCCGTGAGG
ACGAAACGCCCTCGACGCAT-3' (SEQ ID NO:1),
5'-CTAGATGCGTCGAGGGCGTTTCGTCCTCACGGACTC
ATCAGTGCTGGAGATCGCC-3' (SEQ ID NO:2) (sense and antisense,
respectively). The resulting duplex was ligated into the EcoRI and SpeI sites of pZeoU1EcoSpe to create pU1FIB. All ligation junctions were sequenced to verify the identity and orientation of the insert.

- 24 -

In that the structure described for naturally occurring antisense RNAs is highly similar to that for small nuclear RNAs (snRNAs), essential components of the spliceosome complex that are abundant and stable in the nucleus of mammalian cells, the U1 snRNA gene was selected as the framework for vector construction (Guthrie, et al., *Annu. Rev. Genet.*, 22; 387, 1988). Other attributes include the potent and constitutively active nature of the U1 snRNA promoter, the ability of the unusual trimethylguanosine 5' cap and Sm protein interactions to signal transport of U1 snRNA back into the nucleus (Hamm, et al., *Cell*, 62; 569, 1990; Fischer, et al., *Science* 249; 786, 1990; Fischer, et al., *J. Cell Biol.* 113 Mo; 705, 1991; Plessel, et al., *Mol. Cell. Biol.* 14, 4160, 1994), and the lack of polyadenylation of mature snRNAs, a factor which may favorably influence transcript trafficking and localization (Zhong, et al., *Proc. Natl. Acad. Sci. USA*, 91; 4258, 1994). Moreover, unlike other spliceosome components, U1 snRNA is widely dispersed in the nucleoplasm (Carmo-Fonseca, et al., *EMBO J.*, 10; 195, 1991; Matera, et al., *J. Cell. Biol.*, 121; 715, 1993).

As an illustrative example, the antisense targeting 'core' was designed to contain sequences exactly complementary to coding nucleotides 1-15 and 17-30 of fibrillin mRNA (Corson, et al., *Genomics*, 17; 476, 1993; Pereira, et al., *Hum. Mol. Genet.*, 2; 961, 1993) separated by the 22 bp hammerhead ribozyme loop. Dominant-negative forms of fibrillin-1 cause Marfan syndrome, an autosomal dominant systemic disorder of connective tissue (Dietz, et al., *Nature*, 352; 337, 1991). The regions of complementarity were predicted to align the autocatalytic structure with the consensus sequence for ribozyme cleavage (5'-GUC-3') within the target message (Cech, et al., *Annu. Rev. Biochem.*, 55; 599, 1986; T.R. Cech, *Science*, 236; 1532, 1987; T.R. Cech, *Ann. Rev. Biochem.*, 59, 543, 1990; Haseloff, et al., *Nature*, 334; 585, 1988). This antisense targeting sequence was substituted for the short Sm protein binding sequence between the two hairpin loops of U1 snRNA (Guthrie, *Annu. Rev. Genet.*, 22; 387, 1988.) The sequence of the resulting chimeric RNA (Figure 1) was analyzed using a program that predicts RNA structure (Mike Zuker's RNA page: <http://ibc.wustl.edu/~zucker/rna/>). The boundaries of the targeting sequence utilized in the construct were selected to maximize preservation of the U1

- 25 -

snRNA stem-loops, the ribozyme secondary structure and the accessibility of the sequence complementary to the target message. Expression of the chimeric RNA was under the transcriptional control of a region of U1 snRNA 5'-flanking sequence that has been shown to be potent and constitutively active (Zhuang, et al., *Cell*, 46; 827, 5 1986; Maugin, et al., *EMBO J.*, 5; 987, 1986); and Asselbeyers, et al., *Mol. Biol. Rep.* 17; 101, 1993).

Figure 2 shows the sequence and predicted structure of chimeric transcripts derived from pU1/FIB. The Sm protein binding site of U1 snRNA (boxed) has been substituted by an antisense targeting core (flanked by dashed lines) that is 10 complementary to the first 30 coding nucleotides of FBN1 and contains a hammerhead ribozyme sequence in its center. The two stem-loop structures of U1 snRNA are maintained. Cleavage of the target message (unbolded characters) is predicted to occur immediately following the consensus sequence 5'-GUC-3' (arrow). The 5' cap structure of the chimeric targeting molecule may contain 1 or 3 methyl (m) 15 groups. A sequence in the chimeric molecule (5'-AAUUGG-3', underlined) remains highly similar to the consensus for Sm protein binding, PuA(U)nGPu (Pu = A or G).

A human osteosarcoma (MG63) cell line was stably transfected with either a reporter gene expression vector (pZeoSVLacZ, Invitrogen) or the chimeric construct (pU1/FIB) by long-term selection for zeomycin resistance. Human osteosarcoma 20 (MG63) cells (Sakai, et al., *J. Cell Biol.*, 103; 2499, 1986) were grown to 60% confluency and transfected with either linearized pU1/FIB or a lac Z reporter gene construct (pZeoSVLacZ, Invitrogen), used as a control. The transfections were performed using a DOTAP liposome formulation (Boehringer Mannheim, 1 mg/ml) according to manufacturer's instructions. Cells were grown in MEM media (Cellgro) 25 with 10% FCS and 250 µg/ml of zeocin (Invitrogen), and were maintained in selection for 14 days prior to the isolation of clonal colonies. Mono- and polyclonal colonies were established and grown to confluency.

Figure 3 shows an immunohistochemical analysis of MG63 cells that were stably transfected with an expression construct (pZeoSVLacZ, Invitrogen) lacking 30 targeting sequence (panels A and C) or pU1/FIB (panels B and D). The antibodies

- 26 -

either recognized epitopes in fibrillin-1 (A and B) or fibronectin (C and D). Analysis was performed in duplicate for each of three independent clonal colonies for each construct, with identical results. Representative fields are shown at 63X magnification. Untransfected MG63 cells showed a pattern of protein deposition that

5 was indistinguishable from that for cells transfected with pZeoSVLacZ.

Fibrillin-1 was strikingly absent upon immunohistochemical analysis (Eldadah et al., *J. Clin. Invest.*, 95; 874, 1995) using either anti-fibrillin-1 mAb 69 (a gift from L. Y. Sakai) or an anti-fibronectin mAb (Sigma) of cells harboring pU1/FIB, while cells transfected with pZeoSVLacZ showed a pattern of protein

10 deposition that was indistinguishable from untransfected controls (Figure 3). The lines harboring pU1/FIB could not be distinguished from mock-transfected or untransfected cells upon immunohistochemical analysis with a monoclonal antibody to fibronectin, suggesting specificity for the targeting process.

Antisense RNAs expressed by pU1/FIB might inhibit fibrillin-1 expression

15 by multiple mechanisms. Ribozyme cleavage would remove the 5' cap from targeted transcripts, an event predicted to effect their rapid degradation. Alternatively, in the absence of ribozyme cleavage, involvement of the initiating AUG codon in duplex formation would impair translation.

The magnetic porous glass direct mRNA purification technique was used

20 to isolate poly(A) RNA (according to the manufacturer's instructions, CPG, Inc.) and Northern blot analyses were performed. Electrophoresis of 3.5 µg of mRNA was performed under denaturing conditions, as previously described (Lehrach, et al., *Biochemistry*, 16; 4743, 1977; D.A. Goldberg, *Proc. Natl. Acad. Sci. USA*, 77; 5794, 1980). The gel was exposed to 60 mJoules UV light to facilitate transfer of large

25 mRNA species. RNA was transferred to nylon membrane using the turboblotting system according to manufacturer's instructions (Schleicher & Schuell). The membrane was washed briefly in 2X SSC, crosslinked with 125 mJoules of UV light and prehybridized in Expresshyb (Clontech) for 1/2 hour at 68° C. Human cDNA probes encoding fibrillin-1 (nt 370-1183), β-actin (Clontech), and G3PDH (Clontech)

30 were labeled by random priming (Feinberg, et al., *Anal. Biochem.*, 132; 6, 1983). The

- 27 -

membrane was hybridized for 1 hour at 68° C, washed first in 2X SSC with 0.05% SDS at room temperature and then in 0.1X SSC with 0.1% SDS at 50° C.

Figure 4 shows a Northern blot analysis of 3.5 µg of poly(a) RNA extracted from confluent MG63 cells that were untransfected (lane 1), or stably
5 transfected with either pZeoSVLacZ (Lane 2) or pU1/FIB (lane 3-5). Cells for lanes 1 and 3 were polyclonal, while cells for lanes 2, 4 and 5 were derived from monoclonal colonies. Hybridization was performed with radiolabeled cDNA probes complementary to fibrillin-1, β-actin, and G3PDH transcripts.

Northern blot analysis of multiple clonal colonies harboring pU1/FIB
10 revealed undetectable levels of FBN1 message (Figure 4). FBN1 transcripts were easily detected upon Northern analysis of RNA extracted from untransfected and mock-transfected cells. All cell types showed comparable amounts of β-actin and G3PDH RNA, confirming the specificity of targeting.

A similar U1 construct was created containing U1 snRNA stemloop
15 structures and an antisense core aimed at inhibiting rent-1 expression. Functional analysis of transfected cells showed a significant up-regulation of the steady-state abundance of transcripts derived from endogenous nonsense alleles.

SUMMARY

These data suggest that most, if not all of the inhibitory effect was
20 achieved at the level of target message abundance, presumably by reducing mRNA stability. Although an influence upon transcriptional efficiency cannot be excluded, it is not intuitive given the site and nature of the predicted targeting sequence-target interaction. Based upon previous studies, the antisense-induced mRNA degradation most likely takes place within the nucleus (Murray, et al., in *Modern Cell Biology*,
25 J.A.H. Murray, Ed. (Wiley/Liss, New York, 1992), pp 1-49 M.Cornelissen, *Nucleic Acids Res.*, 17; 7203, 1989.). Three elements of nuclear pre-U1 RNA, the 5' m7G cap, the 3' terminal stem loop, and sequences in the 5'-terminal 124 nucleotides, contribute to the stability of pre-U1 and snRNA and to efficient nuclear export (Neuman de Vegvar, et al., *Mol. Cell. Biol.*, 10; 3365, 1990; Terns, et al., *Genes Dev.*,

- 28 -

- 2; 1898, 1993; Yuo, et al., *ibid.*, 3; 697, 1989; Izaurralde, et al., *Nature*, 376; 709, 1995). Nuclear targeting of cytoplasmic U1 snRNA is influenced by hypermethylation of the 5' cap structure and binding of at least one common U snRNP protein, both dependent upon the integrity of the Sm protein binding site which has
- 5 been altered in pU1/FIB (Guthrie, et al., *Annu. Rev. Genet.*, 22; 387, 1988). A sequence at the 5' end of the insert (AAUUGG) is highly similar to the consensus site [PuA(U)nGPu, Pu = A or G] for Sm protein binding (Jones, et al., *EMBO J.*, 9; 2555, 1990.). It has also been shown that Sm binding sites are highly tolerant of mutations including internal nucleotide substitutions and deletions (Jones, et al., *supra*).
- 10 Stable transfection of cultured cells with an expression vector containing the isolated FBN1/ribozyme core sequence of pU1/FIB but no U1 snRNA secondary structure, under the transcriptional control of the SV40 promoter, resulted in no discernible inhibitory effect on the expression of fibrillin-1 transcript or protein. The oligonucleotides described above containing the 24 highly conserved nucleotides of
- 15 hammerhead ribozymes flanked by complementary sequence for 10 codons in the translation start site region of the FBN1 mRNA, were cloned into the SpeI and EcoRI restriction sites for the polylinker of pZeoSV (Invitrogen). Fetal fibroblasts were stably transfected with this vector. Immunohistochemical analysis of these clones was performed by methods identical to those described herein). These results suggest
- 20 that selected properties of U1 snRNA and/or its promoter are contributing to the extreme efficiency of inhibition seen with the use of the chimeric cRNA.

Although the invention has been described with reference to the presently preferred embodiment, it should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention

25 is limited only by the following claims.

- 29 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: The Johns Hopkins University School of Medicine
- (ii) TITLE OF INVENTION: Delivery Construct for Antisense Nucleic Acids and Methods of Use
- (iii) NUMBER OF SEQUENCES: 2
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Fish & Richardson P.C.
 - (B) STREET: 4225 Executive Square, Suite 1400
 - (C) CITY: La Jolla
 - (D) STATE: CA
 - (E) COUNTRY: USA
 - (F) ZIP: 92037
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/742,943
 - (B) FILING DATE: October 31, 1996
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Haile, Lisa A.
 - (B) REGISTRATION NUMBER: 38,347
 - (C) REFERENCE/DOCKET NUMBER: 07265/098WO1
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 619/678-5070
 - (B) TELEFAX: 619/678-5099

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 55 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)

- 30 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

AATTGGCGAT CTCCAGCACT GATGAGTCCG TGAGGACGAA ACGCCCTCGA CGCAT 55

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 55 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CTAGATGCGT CGAGGGCGTT TCGTCCTCAC GGACTCATCA GTGCTGGAGA TCGCC 55

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A nucleic acid construct for suppressing expression of a target gene, comprising:
 - 5 an antisense nucleic acid sequence directed to a target gene of interest;
 - 5' of said antisense nucleic acid region, an unmodified, natural occurring 5' U snRNA stem loop structure;
 - a pol II promoter region 5' of said antisense nucleic acid sequence;
 - 10 and
 - 3' of said antisense region, an unmodified, naturally occurring 3' U snRNA stem loop structure,
 - wherein said expression of said target gene is suppressed by at least 75% of the normal level of expression.
- 15 2. The nucleic acid construct of claim 1, wherein the U snRNA is U1.
3. The nucleic acid construct of claim 1 or claim 2 wherein the promoter is a U1 snRNA promoter.
- 20 4. The nucleic acid construct of claim 1 or claim 2 wherein the promoter is a constitutive promoter.
5. The nucleic acid construct of claim 1 or claim 2 wherein the promoter is an inducible promoter.
- 25 6. The nucleic acid construct according to any one of claims 1 to 5, further comprising a ribozyme nucleic acid.
- 30 7. The nucleic acid construct of claim 6, wherein the ribozyme nucleic acid is located between the 5' and 3' stem loop structures.
8. The nucleic acid construct of claim 6 or claim 7 wherein the ribozyme nucleic acid is a hammerhead-type ribozyme.



9. The nucleic acid construct of claim 6 or claim 7 wherein a consensus sequence for ribozyme cleavage in a target nucleic acid is 5'-GUC-3' or 5'-GUA-3'.

5

10. The nucleic acid construct according to any one of claims 1 to 9, wherein the antisense nucleic acid is targeted to a region of a gene is selected from the group consisting of rent-1, HPV E6, HIV, hyaluronic acid synthase, and fibrillin.

10

11. A method for suppression of gene expression comprising administering to a cell a suppressive-effective amount of the nucleic acid construct according to any one of claims 1 to 10.

15

12. The method of claim 11, wherein the administering is *in vitro*.

13. The method of claim 11, wherein the administering is *in vivo*.

14. The method of claim 11, wherein the administering is *ex vivo*.

20

15. The method according to any one of claims 11 to 14, further comprising:
administering a second nucleic acid encoding a wild-type polypeptide corresponding to the gene product of the gene being suppressed, wherein the second nucleic acid is resistant to ribozyme cleavage and/or antisense inhibition.

25

16. A non-human transgenic animal comprising the nucleic acid construct according to any one of claims 1 to 10.

30

17. A non-human knock-out animal produced utilizing the nucleic acid construct according to any one of claims 1 to 10.

18. The nucleic acid construct according to any one of claims 1 to 10, further comprising a 5' trimethylguanosine cap.



Dated this fifteenth day of November 2000

The Johns Hopkins University School
of Medicine

Patent Attorneys for the Applicant:

F B RICE & CO

Patent Office
Australia



FIG. 1





FIG. 2



Plasmid name: pU1/RIBOZYME
Plasmid size: 3610 bp
Constructed by: H. C. Dietz & R. A. Montgomery
Construction date: 11/20/95
Comments/References: none

FIG. 3




FIG. 4

